

Characterization and variability of the prokaryotic community in sediments from Salar de Lagunilla, Northern Chile

Caracterización y variabilidad de la comunidad procariota en sedimentos del Salar de Lagunilla, Norte de Chile

CRISTY A. MEDINA ARMIJO & RUBEN MORAGA MAMANI

Laboratorio de Ecología y Microbiología Ambiental, Facultad de Recursos Naturales Renovables, Universidad Arturo Prat, Av. Arturo Prat 2120, Iquique, Chile
E-mail: rmoraga@unap.cl

ABSTRACT

Fluorescence *in situ* hybridization (FISH) with rRNA-targeted oligonucleotide probes was used to investigate the abundance of prokaryotic community from the sediments of different sites and along a temporary-space scale from Salar de Lagunilla, a high-altitude athalassohaline wetland in the Chilean Altiplano. Six different taxonomic groups were studied: *alpha*, *beta*, *gamma* proteobacterias, sulfate-reducing of the delta subclass of *Proteobacteria*, *Cytophaga-flavobacteria* and the domain *Archaea*. The analyses showed the *Archaea* domain and *Cytophaga-flavobacteria* group (33.6 % - 20.2 %, respectively) as the dominant group, whereas the groups with minor abundance corresponded to *alpha* and *Beta-proteobacteria* (10.7 - 15.5 %). The prokaryotic communities in the sediments also developed differently, as shown by Fluorescence *in situ* hybridization (FISH) and by nonmetric multidimensional scaling analysis. Changes in prokaryotic community composition were followed during three years, where there was a difference in the abundance of the domain *Archaea*, during the dry season (March-August 2006), which demonstrated the existence of temporary differences, but not space. In addition, Canonical correspondence analysis revealed that the prokaryotic community composition could be influenced by some environmental factors, where important components of the ecosystem such as, limestone and carbon, could play a role in the distribution and composition of the prokaryotic community.

KEYWORDS: fluorescence *in situ* hybridization, salt flats, Andean Altiplano.

RESUMEN

Se estudio la composición procariota de los sedimentos asociados a la laguna del salar alto andino de Lagunilla, norte de Chile, mediante la técnica de Hibridación *in situ* Fluorescente (FISH), a lo largo de una escala espacio-temporal. Se estudiaron seis diferentes grupos de la comunidad procariota *alfa*, *beta*, *gamma* protobacterias, *sulfatoreductoras*, *Cytophaga-flavobacterias* y el dominio *Archaea*. Los análisis mostraron que el grupo más dominante fue el Dominio *Archaea* y *Cytophaga-flavobacteria* (33.6% y 20.2%), mientras que los grupos con menor abundancia corresponden *alfa* y *beta*-protobacteria (10.7 y 15.5%). Se analizaron los cambios en la composición de la comunidad microbiana, a través de un análisis de escalamiento multidimensional no métrica. Los cambios en la composición de la comunidad procariotas fueron seguidos durante tres años, donde se observó una diferencia en la abundancia del dominio *Archaea*, durante la estación seca (marzo-agosto de 2006), el cual demostró la existencia de diferencias temporales, pero no espaciales. Los análisis de correspondencia Canónica, revelaron que la composición procarionte fue influenciado por algunos factores ambientales, siendo la caliza y el Carbono componentes de importancias en este ecosistemas, los cuales juegan un rol en la distribución y composición de la comunidad procariota.

PALABRAS CLAVES: Hibridación *in situ* Fluorescente, salares, Altiplano Andino.

INTRODUCTION

Salar de Lagunilla is located in the Chilean altiplano, at 3800 m of altitude, is considered an athalassohaline system, because the composition of water is different from the seawater (Oren 2002). The waters of the

lagoon have a variable salinity and pH neutral where the principal ionic composition corresponding to Na⁺, K⁺ and Cl⁻. Abiotic condition in the altiplano including, their geographical isolation and remoteness, high UV radiation and low dissolved oxygen concentration, strong variation in atmospheric pressure and, low temperature in the night

and high temperature in the daily (Rodríguez-Valera 1988; Ventosa 2006). In addition, the Chilean altiplano is characterized by two very different seasons, a wet period (characterized by the input of storm-water during the months of January and February) and dry period with minimal or low rainwater in the rest of year (Karzulovic & Garda 1979). All these factors are crucial for biota inhabiting these environments, to be adapted to this extreme condition.

The study of microbial diversity and the distribution of microorganisms that inhabit in salt lakes, are crucial since these types of microorganism play a fundamental role in biogeochemical cycles such as carbon and nutrients. As example, the study of the microbial composition in saline environments might present potentially interesting in biotechnological processes as the bioremediation, UV-blockade and new antibiotics production. Also the study of their participation in biogeochemical processes as degradation and remineralization of the organic matter could provide information about the evolution of the Earth (Dundas 1998; Hoehler *et al.* 2001).

In the present study, we describe the composition and abundance of prokaryotic community in the sediment samples

at, five contrasting sites from Salar de Lagunilla using Fluorescent *in Situ* Hybridization (FISH) technique has been widely used in studies of microbial ecology (Fazi *et al.* 2007; Pernthaler & Amann 2005; Glöckner *et al.* 2000; Llobet-Brossa *et al.* 1998; Snair *et al.* 1997; Amann *et al.* 1991), providing an important tool to study the composition of microbial communities. In addition, we also studied the changes in temporary space induced by environmental factors, whit sediment samples taken during three years.

MATERIAL AND METHODS

SITE DESCRIPTION AND SAMPLING

The study was developed in the Salar de Lagunilla, located in the Andes mountains to the coast 155 km north of Chile (19°56'01''S 68°50'54''W). There were established five sites of permanent sampling in the lagoon including from north to south, realizing the compilation of the samples between March 2006 and February 2008. (Figure 1)

The samples were extracted from the superficial sediments of the lagoon by core (10-cm of length and 3 cm width), saved immediately to hermetic bags and fixed directly in 1%




FIGURE 1. Map indicating the location of Salar de Lagunilla and the sampling sites.

FIGURA 1. El mapa indica la localización del Salar de Lagunilla y puntos de muestreo.

formalin-phosphate-buffered saline (PBS) (composed of 0,05 M Na₂HPO₄, 0,85% NaCl [pH 7 in water]). The samples were transported at 4°C and processed immediately upon return to the laboratory then, withing 5 h of sampling. From each sample was performed in surface sediment mix which took 0.5 grams, the formalin-fixed samples wer washed in PBS and centrifuged at 10.000 rpm for 5 minutes and then he supernatant is stored under following conditions in ethanol-PBS (1:1) at 20°C. Sample was diluted and treated by mild sonication for 30 pulses each performed a second and then continues with a series of 5 min of centrifugation at 4,000 rpm. Washed sample was taken 30 µl for be filtered in 0.2 µm pore-size polycarbonate filter, 22 mm in diameter with 1 ml of sterile distilled water, yielding two filtered samples, which were subsequently analyzed by FISH.

CY3-LABELED OLIGONUCLEOTIDE PROBES

The specific probes for recognition of the *Bacteria* domain (ALF1b, BET42a, GAM42a, SBR385, CF319a) and *Archaea* (ARCH915), were used for the specific detection of the object cells. Probes with in order to optimize the hibridization conditions, non-fluorescent competitors probes were used under the same conditions of BET42a and GAM42a probes (Manz *et al.* 1992). The specific oligonucleotide sequences of CY3-labeled probes and the hybridization conditions are summarized in the Table 1.

STAINING AND CELL COUNTS OF SAMPLES

To perform FISH, Filters were quartered with a razor blade, ande the section (6 to 7) were placed face up on the glass slide. Aproximately 20 µl of preheated hibridization solution was placed on each section filter, and 50 ng specific probes

were added in each section of filtere. The hibridization solution contained 20 to 35% (p/v) formamide (depending on the experiment; see table 1), and a mixture (5M NaCL, 1M Tris-HCl [pH 7,2], 10% sodium dodecyl sulfate [SDS]) and incubated at 46 ° C for 2 hours.

Afterwards, the filters sections were incubated in 10 ml of prewarmed washing buffer at 48 ° C for 15 min. The washing solution consisted of (20 mM Tris-HCl (pH 8), 5 mM EDTA, 0.01% SDS and variable concentration of NaCl). A formamide concentration of 35% was used 80 mM and 20% was used 0,225 mM of NaCl. The filter sections were dried on blotting paper, placed back on a glass slide, and covered whith 50 µl of DAPI solution, final concentration, 2 to 5 µg/ml, for 5 min at room temperature in the dark. Then, the filters were gently washed in 2ml of 0,2 µm-filtered destiled water and ethanol, dried on blotting paper, and mounted on glass slides.

The filter sections were inspected whith a fluorescence microscope, equipped whith a 50-W high-pressure mercury bulb and specific filter sets (DAPI [Zeiss 01]), Cy3 (Chroma HQ 41007). Each microscopic field was first viewed with CY3 filter, before switching to the DAPI filter, to avoid bleaching of CY3 during the DAPI count. For each samples and probe, more tan 500 cells were counts; for the DAPI examination, more then 1500 cells were counted per samples. All probe-specific cell counts were presented as the percentage of cell counts by DAPI. The mean abundances and standard deviations were calculated from the counts of 10 to 20 randomly chosen fields on each filter section. The account of the abundance of prokaryotic cells per gram was performed using the protocol and algorithms proposed by (Fry 1986).

TABLE 1. Oligonucleotide Sequences and Percentage of formamide (% FA).

TABLA1. Secuencia de Oligonucleotidos y porcentajes de formamida.

Probe	Secuencia (5'-3')	Posición en ARNr ^a	Specificity	% FA <i>in situ</i> ^b	Reference
ALF 1b	GGTAAGGTTCTGCGCGTT	16S (968-986)	<i>α-proteobacteria</i> ,	20%	Manz <i>et al.</i> 1992
BET 42 ^a	GCCTTCCCACTTCGTTT	23S (1027-1043)	<i>β-proteobacteria</i>	35%	Manz <i>et al.</i> 1992
Gam 42 ^a	GCCTTCCCACATCGTTT	23S (1027-1043)	<i>γ-proteobacteria</i>	35%	Manz <i>et al.</i> 1992
CF319a	TGGTCCGTGTCTCAGTAC	16S (319-336)	<i>Cytophaga-flavobacteria</i>	35%	Manz <i>et al.</i> 1996
SBR 385	CGGCGTCGCTGCGTCAGG	16s (385-402)	<i>δ-protobacteria</i>	35%	Amann <i>et al.</i> (1990a, 1995b)
ARCH915	GTGCTCCCGCAATTCCT	16S (915-934)	<i>Archaea</i>	35%	Stahl & Amann, 1991

^a *Escherichia coli*, numbering (Brosius . 1981).

^b Percent formamide (FA) in situ hybridization buffer.

STATISTICAL ANALYSIS

Of the data matrix was constructed Bray-Curtis similarity. A non-metric multidimensional scaling analysis (NMDS) was performed to study the changes in the distribution patterns and variability of the prokaryotic community. Their similarity analysis comparing the sampling sites were performed by ANOSIM (Analysis of Similarity). In this sense, ANOSIM shows an R-value statistic, which determinate the distance ranges within and between groups or samples. In general this value can be divided into 3 categories; $R > 0,75$: indicates that there are large differences and treatments or groups are well separated, $R > 0,5$: indicates (groups/samples) as separated, but overlapping, and $R < 0,25$: indicates little or no differences, and treatments or groups are barely separable (Clarke 1993). To avoid differences in the distribution of data by the sample size, proportions were transformed to $\log(x + 1)$ (Ramette 2007). All tests were carried out using PRIMER software version 6 (Clarke & Gorley, 2005).

To study if the composition of biological community was influenced by environmental parameters, a canonical correspondence analysis (CCA) was performed using the Community Ecological Analysis software (Ecom II version 2.01),—(Seaby & Henderson, 2007), was used to compared the prokaryotic abundance (FISH) matrices with the environmental variables (physicochemical matrices). Both FISH data and environmental variables were $\log(x + 1)$ transformed. Forward selection was used to rank the environmental variables in importance for determining the species data (Multicollinearity test). The environmental (explanatory) variables used were nitrogen concentration measured as NO_3 , percentage of limestone measured as (CaCO_3) , percentage of easily oxidized carbon (CO) and electrical conductivity (EC [mS/cm]).

RESULTS

ESTIMATING THE ABUNDANCE OF PROKARYOTES

Table 2 shows the variation of cell counts per gram (cell g^{-1}) and percentages of FISH counts during the months of the study. The maximum value of DAPI counts ($6,98 * 10^7 \text{cells g}^{-1}$) was detected during August 2006 and the minimum ($2,39 * 10^7 \text{cells g}^{-1}$) was detected in April 2007; being the average DAPI counta total of $3,26 * 10^7 \text{cells g}^{-1}$. In addition, the percentage of FISH detectable cells to the *Bacteria* and *Archaea* domains was $9,05\text{E} +05$ and $1,08\text{e} +06$ (cells/g) respectively.

These sediments have a high diversity of cell morphologies as, filamentous bacillary forms, bow-shaped rod, coccoid, flagellum cell, bacilli in chains, and among them were large consortia bacteria. Most of the cell identifica in the sediment showed a homogeneous morphology of thin short road. (Figure 2).




FIGURE 2. Abundance procariota of the sediments associated with Salar de Lagunilla in cell/ grams. (A) Alfa-proteobacteria (B) Beta-proteobacteria (C) Gamma-proteobacteria (D) Cytofaga-flavobacteria (E) Bacteria Sulfatoreductoras (F) Archaea.

FIGURA 2. Abundancia procariota de los sedimentos asociados al Salar de Lagunilla en Células/gramos. (A) Alfa-proteobacterias (B) Beta-proteobacterias (C) Gamma-proteobacterias (D) Cytofaga-Flavobacterias (E) Bacterias Sulfatoreductoras (F) Archaea.

The groups with major variability were the domain *Archaea* and *Alpha-proteobacteria* with relative percentages of hybridized cells (FISH) were from 11.8 ± 5.2 to 34.7 ± 10.7 % and from 7.9 ± 2.0 to 31 ± 8.8 % respectively, whereas that the unalterable groups were *Gamma-proteobacteria* and *Cytofaga-flavobacteria* with relative percentages of hybridized cells from 15.7 ± 4 to 22.5 ± 8.5 % and from 14.6 ± 10 to 28.7 ± 2.3 % respectively. On the other hand, the most abundant group corresponded to the *Archaea* domain with relative abundances of 34.7 % in April, and 29% during the months of March and August (2006), and from November to February (2007). In the same way, the less abundant detected group was *Beta-proteobacteria*, showing a relative abundance of 5.5% during March 2006. Members of *sulfate-reducing* bacteria group from the subclass *Delta-proteobacteria*, showed an increase in their relatives abundances during the months of November, 2007 and February, 2008 (25.0 ± 2.5 and 27.5 ± 1.2 %, respectively), whereas the minor relatives abundances were observed during the months of November, January and April, 2007.

In general the sediments system of the lagoon showed a trend where the domain *Archaea* and the group *Cytofaga-flavobacteria* were dominant. The members of the *sulfate-reducing* and *Gamma-proteobacterias* presented similar abundances. *Alpha-proteobacterias* and *Beta-*

proteobacteria group were the group with less abundances in the superficial sediments system of the salar de Lagunilla.

PHYSICOCHEMICAL RELATIONSHIP OF THE PROKARYOTIC COMMUNITY.

To describe the relations between the abundances of the prokaryotic community and the physicochemical variables (Table 3), there was made a canonical correspondence analysis (CCA). The results for the humid period (March, 2006, January, 2007 and February, 2008), makes clear 48.7 % of the accumulative variance, on the other hand the dry period (August, 2006, November, 2006, April, 2007, August, 2007 and November, 2007), the variable significant for the first two axes explain 45.2 % of the accumulative variance.

The table 4 shows the environmental variables, which contribute, significantly in the Dry period, from bigger to minor contribution in model they are, NO_3 mg/K, Limy (CaCO_3 g), % p/p CO (carbon easily oxidabl) and CE mS/cm. For the humid period the significant concentrations were: % p/p CO, NO_3 mg/k, limestone (CaCO_3 g), CE mS/cm and PO_4^{+3} mg/k expressed in the axis 1.

Biplot diagram ordination of CCA analysis shows the patterns of changes in prokaryotic community, which can be explained by environmental variables.

TABLE 2. Total DAPI count and specific FISH studies for months.

TABLA 2. Recuento total DAPI y FISH por meses de estudios.

Month of Study	Total cell counts (cel/g) (Mean)	Fraction (%) of total cell (mean \pm SD) detected with probea:					
		ALF968	BET42a	GAM42a	CF319a	Sbr385	Arch915
mar-06	2.90E+07	7.9 \pm 2.0	5.5 \pm 0.5	17.0 \pm 3.9	NA _b	17.4 \pm 0.9	34.4 \pm 3.8
aug-06	6.08E+07	11.1 \pm 1.1	17.0 \pm 0.7	18.5 \pm 3.7	NA _b	18.5 \pm 1	33.5 \pm 2.8
nov-06	2.95E+07	19.0 \pm 6.9	16.9 \pm 4.5	20.4 \pm 10.9	14.6 \pm 10	13.1 \pm 12.6	13.6 \pm 7.2
jan-07	2.39E+07	14.6 \pm 5.7	21.2 \pm 8.3	15.7 \pm 4	21.3 \pm 8.1	13.7 \pm 4.8	19.2 \pm 3.2
apr-07	2,42E+07	12.3 \pm 4	25.1 \pm 5.1	21.0 \pm 8.1	20.8 \pm 3.7	14.8 \pm 4.1	34.7 \pm 10.7
aug-07	2.84E+07	12.6 \pm 5.6	14.8 \pm 3	18.8 \pm 6.8	18.8 \pm 3.4	19.5 \pm 7.8	11.8 \pm 5.2
nov-07	2.76E+07	31.0 \pm 8.8	26.7 \pm 5.9	22.5 \pm 8.5	23.0 \pm 5.1	25.0 \pm 2.5	29.6 \pm 1
feb-08	2.73E+07	19.0 \pm 4.5	20.6 \pm 4	17.6 \pm 3.4	28.7 \pm 2.3	27.5 \pm 1.2	28.7 \pm 1

a Detection rate compared with DAPI. Mean and standard deviation were calculated by counting 20 fields in the section of the filter.

b Unparsed

TABLA 3. Propiedades Físico-Químicas del Salar de Lagunilla.

TABLE 3. Physical-Chemical Properties of the Salar de Lagunilla.

Month of Study	pH	CE (mS/cm)	%CO	% Sulfato	% Caliza (CaCO ₃)	NO ₃ (mg/K)	PO ₄ ⁺³ (mg/K)
March-2006	8.1	9.0	2.1	1.5	16.4	ND ^a	ND ^a
August-2006	7.5	6.3	1.6	1.1	13.7	ND ^a	ND ^a
November-2006	7.5	8.0	2.1	0.9	14.0	ND ^a	ND ^a
January -2007	7.5	8.5	1.7	0.9	13.8	24.4	7.1
April -2007	7.9	4.8	1.8	0.9	13.0	28.7	9.3
August-2006	8.3	4.1	1.4	0.8	15.0	25.5	7.9
November-2006	7.1	1.7	1.4	0.6	15.1	22.7	9.6
February -2008	7.8	8.5	1.4	0.8	ND	18.9	11.2

ND^a, a undetermined

TABLA 4. Intranet values of environmental variables

TABLE 4. Valores intranet de las variables ambientales

Environmental variables		1	2
dry period	CE (mS/cm)	0.04	0.91
	% p/p CO	0.56	0.20
	CaCO ₃ (g)	0.81	0.15
	NO ₃ (mg/k)	0.90	0.09
Wet Period	CE (mS/cm)	0.14	0.40
	% p/p CO	0.57	0.12
	CaCO ₃ (g)	0.24	0.72
	NO ₃ (mg/k)	0.48	0.45
	PO ₄ ³ (mg/k)	0.09	0.05

The environmental variables are represented by the vectors, which it is constructed by the points of the variables on the maximum inertia projected in the space. The arrows represent the vector direction. The vector length indicates the importance of the variable in the model, and the position of these indicates that so correlated the environmental variable is with the axes. Also the angles between the vectors indicate the correlation that exists between them (a small angle indicates high correlation). The samples are

represented by the pictures and indicate the relation with the environmental variables (vector) and his arrangement with environmental variables in the model.

The figure 3 shows the analysis corresponding to the dry period, for this period the available NO₃ and the Limestone (CaCO₃) they are the most important variables for the model, since 1 is narrowly tied to the axis.

The group *Cytofaga-flavobacteria* shows a good correlation of negative trend with the limy variable (CaCO_3) and *Gamma-protobacteria* correlates negatively with the electrical conductivity (CE), explained in the axis 2, this one can represent important quantities of you work out soluble.

For the humid period (Figure 3), the most influential variables in the model are the organic carbon easily to oxidize (CO) and the available NO_3 . The members of the *Sulfato-reductors* it is correlated by the NO_3 positively, *Beta-protobacterias* is correlated negatively with CaCO_3 , which explains to itself in the axis 2, *Gamma-protobacteria* with organic carbon easily oxidable (CO), *Alpha-protobacteria* and the Domain *Archaea* show correlations positive with electrical conductivity, which explains to itself in the axis 2 and *Cytofaga-flavobacteria* is not correlated by any variable, in general the model shows a correlation with the samples and the variable electrical conductivity (CE).

SPATIO-TEMPORAL VARIABILITY

In term of the relative abundances of the prokaryotic

community, these exhibited a great variability during the study period. In the figure 1 NMDS, shown that March and August, 2006 are different from the rest of the study, these are characterized by presenting high abundances of microorganism of *Archaea* domain (34.4 % and 33.5 % respectively), whereas the sum of the relative abundances between the groups of the domain *Bacteria* represent 47.8 % and 65,1 % respectively, with regard to the rest of the groups analyzed in comparison to other studies periods. On the other hand the figure 4 shows the analysis spreads of NMDS, in this analysis one does not show significant differences between the stations of sampling. For this study there were identified 5 sites of study (L1-L2-L3-L4-L5).

The abundance of the prokaryotic community was compared to be able to observe differences intra and interannual. The ANOSIM showed significant differences with overlapping during the period of study ($R = 0.527$, $p < 0.001$), also an ANOSIM was realized to observe the spatial differences, where significant differences were not observed ($R = 0.102$, $p < 0.005$).




FIGURE 3. Canonical correspondence analysis of the prokaryotic abundance and the environmental variables. (A) Humid period and (B) Dry period.

FIGURA 3. Análisis de correspondencia canónica de la abundancia procariotas y las variables ambientales. (A) Período Húmedo y (B) período seco.




FIGURE 4. (A) NMDS, temporary scale representation of the community prokaryotic abundances. Log transformation (X+1), Stress=0.11 and Randon 999 permutations. (B) NMDS, spatial scale representation of the community prokaryotic abundances. Log transformation (X+1), Stress = 0.18 and Randon 999 permutations.

FIGURA 4. (A) NMDS Representación a escala temporal de las abundancias procariotas comunidad. Transformación Log (X + 1), Estrés = 0.11 y Randon 999 permutaciones. (B) NMDS, representación espacial escala de las abundancias procariotas comunidad. Transformación Log (X + 1), Estrés = 0,18 y Randon 999 permutaciones.

DISCUSSION

The Highland Lakes as the Salar de lagunilla represent unique and extreme habitats clearly dominated by various forms of microbial *life*. However, microbiological studies have been performed only occasionally (Demergaso *et al.* 2004; Demergaso *et al.* 2008; Dorador *et al.* 2008a, 2008b, 2009, 2010). Salar de Lagunilla is considered as a moderately athalassohaline wetland (Table 3), comparable in its phylo-genetic groups with other high-altitude cold aquatic systems (Tibetan Lakes: Dong *et al.* 2006; Jiang *et al.* 2006. Atacama Lakes: Demergasso *et al.* 2004)

In this work, using FISH technique was possible to establish a first approximation of the microbial composition and the most abundant specific groups in the sediments of the Salar de Lagunilla. However, the use of this technique has its

limitations, such as the weak signal and lack of specificity of the oligonucleotides. These limitations have been described and solutions have been proposed by Penthaler & Amann, (2005); thus been possible to be more objective with our results. Moreover it is important to note that the contribution achieved through these studies will help to generate other future investigations.

Bacterial assemblages were dominated by *Cytofaga-Flavobacteria-Bacteriodes* (CFB), (28.7% abundances of hybridized cells). This percentage is high when considering the percentage of coverage with the probe, which is about 38% (Amann *et al.* 2008). This group is characteristic in the oceans where it is abundant between the groups (Kirchman 2002), just as this group has been reported in sediments and in environments with high concentrations of salinity (Antón *et al.* 1999). These results are consistent with others studies,

where has been observed in athalossahalino lake of the Atacama by PCR-denaturing gradient gel electrophoresis (DGGE) and sequencing of 16S rRNA gene fragments where the tendency for increasing contribution of (CFB) for higher salinities and altituded (Demergasso *et al.* 2004). In addition, in the sediment samples from three evaporitic basins located in the highlands of northern Chile, has observed the high diversity of bacteroides through of 16S rRNA gene clone libraries created with targeted Bacteroidetes-specific primers and separation of specifically amplified gene fragments by denaturing gradient gel electrophoresis (DGGE) (Dorador *et al.* 2008a). Similar results have also been found in high-altitude lakes from the Tibetan Plateau (Wu *et al.* 2006).

Cytofaga-flavobacteria have been described for their role in the degrader of complex molecules and high molecular weight cellulose and, has also been seen in the presence of abundant phytoplankton booms and association with some algae that may result in enhanced algal growth (Grossant 1999).

The increase of abundant CF in the wet period, where organic matter and salinity increases, due to the death of organisms from other trophic levels and increased salinity, and your affinity for CaCO₃ apparently show that their role is similar to their role in the ocean. Therefore, this group could be the responsible for the degradation of high molecular weight compounds in altiplanic lakes.

The Proteobacteria group is characteristic as a cosmopolitan and grouped according to their habitat, creating divisions in the sea, fresh water or soil (Glockner *et al.* 2000; Bowman *et al.* 2003 & Crump *et al.* 2004). *Alpha* and *Beta*-proteobacteria are the group with the lowest rates of hybridization; this is consistent with studies in the ocean (Glöckner *et al.* 1999; Cottrell *et al.* 2000;). These groups are not abundant in saline environments, but it includes mostly the halotolerant bacteria. However this group has been associated with salt deposits, but participating in processes of nitrification (Nold & Zwart 1998).

Gammaproteobacteria group showed variability in relative abundance in the order of 19%, which has been observed in the work of (Cottrell *et al.* 2000). This group is characteristic as oxides in aerobic and anaerobic environments (Ollivier *et al.* 1994), and it has also been shown that members of this group are facultative able to adapt to different environmental conditions. In studies recently, has been observed the present of anoxygenic phototrophic bacteria in the Atacama salar, across application of the functional pufLM genes, was determined a new phylogenetic lineage of phototrophic Gammaproteobacteria (Thiel *et al.* 2011). Possibly, this group have a role important as primary producers in the salar of Lagunilla

Members of the sulfate-reducing bacteria, showed variability in the abundance in the order of a 13 - 27% hybridization of cells in the period of study. This group is important for the degradation of organic matter in sediments, using sulfate as electron acceptor (Ollivier *et al.* 1994). In the salar de Huasco, has been observed the present of deltaproteobacteria, with a study of 16S rRNA gene sequences related to sulfate-reducing bacteria of the genera *Desulfobacterium* (Dorador *et al.* 2009), this group can have a significant fraction of carbon mineralization in sediments hypersaline.

The salar de Lagunilla is considered as an athalassohaline enviroments, also this system is highly variable depending on the groundwater inputs in different zones of the lake basin and stormwater inputs, especially in the months of January and February, and bacteria have to adapt locally to the new conditions. In this studie the abundance of archaeas comunities exhibit high contribution in this system above bacterium. Howeber has been described that dominance of Archaea over Bacteria in enviroments where NaCl concentration close to saturation, here the halobacterias are the dominant (Oren 2002). However, has been observed the predominance of methanogenic archaea and ammonia-oxidizing Archaea in studies in Salar de Huasco (Dorador *et al.*, 2010). Probably, the Salar de Lagunilla presents dominance by methanogenic archaea, because physical-chemical characteristics of the Lagunilla's sediments. This sediment has high carbonates saturation and anoxic sediments forming minerals, being a favorable condition for the adaptability of these organisms.

The environmental variables affected the diversity and composition of bacterial communities strongly. CCA analysis indicated that the salinity gradients, CaCO₃ and carbon organic are the environmental variables more significantly close 50% of the variance, in specie composition bacterial. The enrichment of carbon and CaCO₃, and the salinity gradient clearly increased the abundance of community bacterial in the sediment; however, the concentration of nitrogen (measured as nitrate) and phosphorus (measured as fosfate) do not significantly affect the abundance and bacterial diversity.

The temporal variability in bacterioplankton communities was significantly highly was highly correlated with salinity, %CO (organic carbon) and CaCO₃; In contrast, spatial variability was relatively small in communities across environments altiplanicos, confirming the observations of microbial diversity and abundance based on FISH. Few studies address temporal and spatial variability using different habitats, Kirchman *et al.* (2010) identified a pattern where differences into winter and summer in the Arctic Ocean of the bacterioplankton communities was

minimal compared with spatial variability across their sampling range. Similarly, Fortunato *et al.* 2012, compares the bacterioplankton community composition from river to surface ocean, by depth from surface to deep ocean, and through time seasonally over an year, by the use of 16S ARN, where they observed changes in the spatial variability and correlations with salinity and Depth.

Our study, show temporary changes that may be associated with increased salinity in dry periods and increased rainfall (wet period), which brings an eutrophication. Although, it seems that the salinity gradient is the best indicator of biogeographic distribution in these environments. We also identified significant overlap in communities of *Archaea* Domain in the dry period.

This study reveals the first analyses of the composition of a prokaryotic community, temporary differences are demonstrated and as the environmental factors they are incidental in the structure, distribution and microbial variability of the populations in the Salar de Lagunilla. These results can serve for future investigations, so that the community analyses could be combined by the evolution of the substrates of the sediments.

BIBLIOGRAPHY

- AMANN, R., SPRINGER, N., LUDWIG, W., GERTZ, H. & SCHLEIFER, K. 1991. Identification *in situ* and phylogeny of uncultured bacteria endosymbiots. *Nature* 351:161-164.
- AMANN, R., LUDWIG, W. & SCHLEIFER, K. 1995. Phylogenetic identification and *in situ*. Detection of individual Microbial Cell without Cultivation. *Microbial Reviews* 59:143-169.
- ANTON, J., LLOBET-BROSSA, E., RODRÍGUEZ-VARELA, F. & AMANN, R. 1999. Fluorescence *in situ* hybridization analysis of the prokaryotic community inhabiting crystallizer ponds. *Environmental Microbiology* 1:517-523.
- BAK, F., & PFENNIG, N.F. 1991. Sulfate-reducing bacteria in littoral sediment of Lake Constance. *FEMS Microbiology Letters* 85:43-52.
- BRAAK, C. 1986. Canonical correspondence analysis: A new eigenvector technique for multivariate direct gradient analysis. *Ecology* 67:1167-1179.
- BOWMAN, J.P., MCCAMMON, S.A., REA, S.M. & MCMEEKIN, T.A. 2000. The microbial composition of three limnologically disparate hypersaline Antarctic lakes. *FEMS Microbiology Letters* 183:81-88.
- BROSIOUS, J., DULL, T.J., SLEETER, D.D. & NOLLER, H.T. 1981. Gene organization and primary structure of a ribosomal RNA operon from *Escherichia coli*. *Journal of Molecular Biology* 148:107-127.
- CLARKE, K.R. 1993 Non-parametric multivariate analyses of changes in community structure. *Australian Journal of Ecology* 18:117-143.
- CLARKE, K.R.; GORLEY R.N. 2006 PRIMER v6: User manual/tutorial, PRIMER-E, Plymouth UK.
- COTTRELL, M.T. & KIRCHMAN, D.L. 2000. Community composition of marine bacterioplankton determined by 16S rRNA gene clone libraries and fluorescens *in situ* hybridization. *Applied and Environmental Microbiology* 66:5116-5122.
- CRUMP, B.C., HOPKINSON, C.S., SOGIN M.L. & HOBIE J.E. 2004. Microbial biogeography along an estuarine salinity gradient: combined influences of bacterial growth and residence time. *Applied and Environmental Microbiology* 70:1494-1505.
- DEMERGASSO, C., CASAMAYOR, E.O., CHONG, G., GALLEGUILLOS, P., ESCUDERO, L., PEDROS-ALIO, C. 2004. Distribution of prokaryotic genetic diversity in athalassohaline lakes of the Atacama Desert, northern Chile. *FEMS Microbiology Ecology* 48:57-69.
- DEMERGASSO, C., ESCUDERO, L., CASAMAYOR, E.O., CHONG, G., BALAGUÍ V. & PEDROS-ALIO, C. 2008. Novelty and spatio-temporal heterogeneity in the bacteria diversity of hypersaline lake tebenquiche (Salar de Atacama). *Extremophiles* 12:491-504.
- DONG, H., G. ZHANG, H. JIANG, B. YU, L. R. CHAPMAN, C. R. LUCAS, & FIELDS, M. W. 2006. Microbial diversity in sediments of saline Qinghai Lake, China: Linking geochemical controls to microbial ecology. *Microbial Ecology* 51:65-82.
- DORADOR, C., BUSEKOW, A., VILA, I., IMHOFF, J. & WITZEL, K.P. 2008A. Molecular analysis of enrichment cultures of ammonia oxidizers from the Salar de Huasco, a high altitude saline wetland in northern Chile. *Extremophiles* 12:405-414.
- DORADOR, C., VILA, I., IMHOFF, J. & WITZEL, K.P. 2008B. Cyanobacterial diversity in Salar de Huasco, a high altitude saline wetland in northern Chile: an example of geographical dispersion?. *FEMS Microbiology Ecology* 64:419-432.
- DORADOR, C., MENESES, D., URRUTIA, V., DEMERGASSO, C., VILA, I., WITZEL, K.P. & IMHOFF I. 2009. Diversity of Bacteroidetes in high-altitude saline evaporitic basins in northern Chile. *Journal of Geophysical Research* 114: G00D05. DOI:10.1029/2008JG000837.
- DORADOR, C., VILA, I., REMONSELLEZ, F., IMHOFF, I. & WITZEL, K.P. 2010. Unique cluster of Archaea in Salar de Huasco, an athalassohaline evaporitic basin of the Chilean Altiplano. *FEMS Microbiology Ecology* 73:291-302.
- DUNDAS, I. 1998. Was the environment for primordial life hypersaline? *Extremophiles* 2:375-377.
- FAZI, S., AMALFITANO, S., PIZZETTI, I. & PERNTHALE, J. 2007. Efficiency of fluorescence *in situ* hybridization for bacterial cell identification in temporary river sediments with contrasting water content. *Systematic and Applied Microbiology* 30:463-470.
- FRY, J. 1988. Determination of biomass. In: *Methods in aquatic bacteriology* (Ed. Austin, B.), pp. 27-72. John Wiley and Sons.
- FORTUNATO C., HERFOR L., ZUBER P., BAPTISTA A., CRUMP B. 2012. Spatial variability overwhelms seasonal patterns in bacterioplankton communities across a river to ocean gradient. *ISME Journal* 6:554-563.
- GLÖCKNER, F., AMANN, R., ALFREIDER, A., PENTHALER, J., PSENNER, R., TREBESIOUS, K. & SCHLEIFER, K.-H. 1996. An *In Situ* Hybridization Protocol for Detection and Identification of Planktonic Bacteria. *Systematic and Applied Microbiology* 19:403-406.

- GLÖCKNER, F.O., FUCHS, B.M. & AMANN, R. 1999. Bacterioplankton compositions of lakes and oceans: a first comparison based on fluorescence *in situ* hybridization. *Applied and Environmental Microbiology* 65:3721-6.
- GLÖCKNER, F.O., ZAICHIKOV, E., BELKOVA, N., DENISOVA, L., PERNTHALER, J., PERNTHALER, A., AMANN, R. 2000. Comparative 16S rRNA analysis of lake bacterioplankton reveals globally distributed phylogenetic clusters including an abundant group of actinobacteria. *Applied and Environmental Microbiology* 66:5053-5065.
- GROSSART, H. P. 1999. Interactions between marine bacteria and axenic diatoms (*Cylindrotheca fusiformis*, *Nitzschia laevis*, and *Thalassiosira weissflogii*) incubated under various conditions in the lab. *Aquatic Microbial Ecology* 19:1-11.
- JIANG, H., DONG, H., ZHANG, H., YU, H., CHAPMAN, L., & FIELDS, M. 2006. Microbial diversity in water and sediment of Lake Chaka an athalassohaline lake in northwestern China, *Applied and Environmental Microbiology* 72:3832-3845.
- KARZULOVIC, J. & GARDA, F. 1979. Evaluación recursos hídricos Provincia de Iquique I Región. Ministerio de Obras Públicas, Dirección General de Aguas; Intendencia I Región; Servicio de Planificación y Coordinación Iquique. Santiago. 205 pp.+ mapas.
- KIRCHMAN, D.L. 2002. The ecology of Cytophaga-flavobacteria in aquatic environments. *Microbial Ecology* 39:91-100.
- KIRCHMAN, D.L., COTTRELL, M.T., LOVEJOY, C. 2010. The structure of bacterial communities in the western Arctic Ocean as revealed by pyrosequencing of 16S rRNA genes. *Environmental Microbiology* 12:1132-1143.
- LLOBET-BROSSA, E., ROSELLÓ-MORA, R. & AMANN, R. 1998. Microbial Community Composition of Wadden Sea Sediments as Revealed by Fluorescence *In Situ* Hybridization. *Applied and Environmental Microbiology* 64:2691-2696.
- MANZ, W., AMANN, R., LUDWIG, W., WAGNER, M. & SCHLEIFER, K.H. 1992. Phylogenetic oligodeoxynucleotide probes for the major subclasses of Proteobacteria: problems and solutions. *Systematic and Applied Microbiology* 15:593-600.
- MATURRANO, L., SANTOS, F., ROSELLÓ-MORA, R. & ANTÓN, J. 2006. Microbial Diversity in Maras Salterns, a Hypersaline Environment in the Peruvian Andes. *Applied and Environmental Microbiology* 72:3887-3895.
- MORIN, P.J. & MCGRADY-STEED, J. 2004. Biodiversity and ecosystem functioning in aquatic microbial systems: a new analysis of temporal variation and species richness-predictability relations. *Oikos* 104:458-466.
- MUYZER, G., WAAL, E.C. & GUITTERLINDEN, A. 1993. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Applied and Environmental Microbiology* 59:695-700.
- NOLD, S.C. & ZWART, G. 1998. Patterns and governing forces in aquatic microbial communities. *Aquatic Ecology* 32:17-35.
- OLLIVIER, B., CAUMETTE, P., GARCIA, J. & MAH, R. 1994. Anaerobic Bacteria from Hypersaline Environments. *Microbiological Reviews* 58:27-38.
- OREN, A. 2002. Molecular ecology of extremely halophilic Archaea and Bacteria. *FEMS Microbiology Ecology* 39:1-7.
- PERNTHALER, J., GLÖCKNER, F., SCHÖNHUBER, W. & AMANN, R. 2001. Fluorescence *in situ* Hybridization (FISH) with rRNA-targeted oligonucleotide probes. In: *Methods in Microbiology: Marine Microbiology* (Ed. Paul, J.H.), pp. 207-226. St. Petersburg, Florida, USA: Elsevier.
- PERNTHALER, J. & AMANN, R. 2005. Fate of Heterotrophic Microbes in Pelagic Habitats: Focus on Populations. *Microbiology and Molecular Biology Reviews* 69:440-461.
- RODRÍGUEZ-VALERA, F. 1988. Characteristics and microbial ecology of hypersaline environments. In: *Halophilic Bacteria* (Ed. Rodríguez-Valera, F.), pp. 3-30. CRC Press, Boca-Raton, Fla.
- RAMETTE, A. 2007. Multivariate analysis in microbial ecology. *Microbial Ecology* 62:142-160.
- RISACHER, F., ALONSO, H. & SALAZAR, C. 2003. The origin of brines and salts in Chilean salars: a hydrochemical review. *Earth-Science Reviews* 63:249-293.
- SEABY, R., HENDERSON, P., 2007. Community Analysis Package 4.0: searching for structure in community data. PISCES Conservation Ltd. Retrieved 09/01/2012 RLU: [http://www.piscs-conservation.com/index.html?softcap.html\\$softwaremenu.html](http://www.piscs-conservation.com/index.html?softcap.html$softwaremenu.html).
- SNAIRD, J., AMANN, R., HUBER, I., LUDWIG, W. & SCHLEIFER, K. 1997. Phylogenetic Analysis and *in Situ* Identification of Bacteria in Activated Sludge. *Applied and Environmental Microbiology* 63:2884-2896.
- SØRENSEN, K.B., CANFIELD, D.E. & OREN, A. 2004. Salinity Responses of Benthic Microbial Communities in a Sola Saltern (Eilat, Israel) *Applied and Environmental Microbiology* 70:1608-1616.
- STAHL, D. A., & AMANN, R. 1991. Development and application of nucleic acid probes. In: *Nucleic acid techniques in bacterial systematics* (Stackebrandt, E. & Goodfellow, M.), p. 205-248. John Wiley & Sons Ltd., Chichester, United Kingdom.
- THIEL, V., TANK, M., NEULINGER, S. C., GEHRMANN, L., DORADOR, C. & IMHOFF, J. F. 2010. Unique communities of anoxygenic phototrophic bacteria in saline lakes of Salar de Atacama (Chile). Evidence for a new phylogenetic lineage of phototrophic Gammaproteobacteria from pufLM gene analyses. *FEMS Microbiology Ecology* 74:510-522.
- VENTOSA, A. 2006. Unusual microorganisms from unusual habitats: hypersaline environments. In: *Prokaryotic diversity-mechanisms and significance* (Eds. Logan, N.A., Lappin-Scott, H.M. & Oyston, P.C.F.), pp. 223-253. Cambridge University Press, Cambridge.
- WU, Q. L., ZWART, G., SCHAUER, M. & HAHN, M. 2006. Bacterioplankton community composition along a salinity gradient of sixteen high-mountain lakes located on the Tibetan Plateau, China. *Applied and Environmental Microbiology* 72: 5478-5485.

Recibido: 19.04.16
Aceptado: 03.11.16